

THE EISENHOWER LIBRARY

3 1151 02695 9787

MORAVIA
SPEC. COLL.

89.406





Wilson

Fioris Daniel

1899

C. A

I. ORTHO-SULPHAMINEBENZOIC ACID.

II. ORTHO-CARBAMINERENZENESULPHONIC ACID.

A Dissertation submitted to the Board of
University Studies of the Johns Hopkins Uni-
versity for the Degree of Doctor of Philosophy.

by

Francis Daniel Wilson.

Baltimore, Maryland.
June, 1899.

-00-

89.456

Contents.

Part I.

Acknowledgement.

Introduction.

Ortho-sulphamine-benzoic Acid, Preparation and

properties of

Salts obtained in Hydrolysis of Benzoic Sulphinide.

Salts in general of Ortho-sulphamine-benzoic Acid, Discussion of the

Neutral Salts of Ortho-sulphamine-benzoic Acid, Preparation and properties of the

Action of Phosphorus Oxychloride upon the Ortho-sulphaminebenzoic Acid.

Action of Ammonium sulphocyanate upon Ortho-sulphaminebenzoic Acid.

Part II.

Introduction.

Ortho-carbaminebenzene-sulphonic Acid, Preparation and Properties of

Salts in general of Ortho-carbaminebenzene-sulphonic Acid.

Neutral Salts of Ortho-carbaminebenzenesulphonic Acid, Preparation and Properties of

Action of Phosphorus Oxychloride upon the Potassium Ortho-carbamine-benzenesulphonate.

40

Action of Ammonia upon the Ortho-carbamine-benzene-sulphone Chloride.

42

Biographical.

Appendix.

Conductivity of Ortho-sulphamine-benzoic Acid and Ortho-carbamine-benzene-sulphonic Acid.

Acknowledgement.

The investigation here presented was suggested by Professor Ira Remsen and was conducted under his personal supervision. I wish, therefore, to express my sincere gratitude for the instruction and inspiration received from him not only in the laboratory but also in the lecture room.

I also wish to acknowledge my gratitude for instruction received from Professor Morse, Professor Clark and Dr. Mathews.

PART I.

ORTHOSULPHUMINEBENZOIC ACID.

Introduction.

Remsen and Fahlberg by oxidizing orthotoluene sul-
phamide with potassium permanganate discovered the sub-
(1)
stance known as benzoic sulphinide. W. A. Noyes by oxi-
dizing ortho-toluene sulphamide with potassium ferricyanide
discovered ortho-sulphamine benzoic acid. He gave to the
acid the formula $C_6H_4 \begin{matrix} \nearrow COOH \\ \searrow SO_2NH_2 \end{matrix}$. The acid is described as
being easily soluble in alcohol and ether and quite readily
soluble in cold water. It crystallized in two different
forms under different conditions: From the barium salt,
acidified with hydrochloric acid and allowed to cool, the
acid separated out in fine needles.

From the barium salt, acidified with hydrochloric
acid and allowed to stand in a cool place for several days,
the acid separated out in large clear prismatic crystals,
which appeared to belong to the monoclinic system.

Mr. Noyes also states that the acid, when heated
rapidly, melts at 165° to 167° . When heated slowly it
melts at 153° - 155° . He thinks this is due to the fact

(1) Ann. Chem., Jour. 8 - 176.

that the acid is partially converted into benzoic sulphide, for if the acid is kept at that temperature for a short while it becomes a solid, and then melts at 210°C . The substance in the tube, he says, has the intensely sweet taste which is so characteristic of benzoic sulphide.

He obtained pure sulphide by heating a portion of the acid to 180°C . in an air-bath. He thinks this change is due to a loss of water of constitution and not of crystallization, as evidenced by the fact that the salts of the acid are totally distinct from the salts of benzoic sulphide.

PREPARATION OF ORTHO-SULPHAMINE-BENZOIC ACID.

Purification of Benzoic Sulphimide.

It occurred to Professor Remsen that the acid could be prepared by boiling pure benzoic sulphimide with sodium hydroxide. The benzoic sulphimide was purified by the (1) method suggested by Pope, which consists in treating the sulphimide with acetone. The sulphimide is more soluble in hot than in cold acetone. The para-sulphamine-benzoic acid, which is sometimes found in benzoic sulphide as an

impurity, is insoluble in acetone. This being the case, it is an easy matter to separate the two substances. The sulphinide used in the following experiments contained scarcely a trace of para-sulphamine benzoic acid, even before it was treated with ~~ether~~ acetone. One hundred grams of the sulphinide required about 460 grams of acetone.

When the benzoic sulphinide is dissolved in hot acetone a white crystalline mass is deposited immediately upon cooling, but if the sulphinide is dissolved in cold acetone and the solution is allowed to evaporate spontaneously massive, colorless rhombohedrons are deposited, some of which are several millimeters in length.

Hydrolysis of Benzoic Sulphinide.

After a number of experiments it was found that when the benzoic sulphinide is treated with sodium hydroxide, under the following conditions, the orthosulphamine benzoic acid was produced in nearly a quantitative yield: Twenty grams of pure benzoic sulphinide was added to a caustic soda solution, which was prepared by dissolving fifteen grams of sodium hydroxide in five hundred c.c. of water. This solution was evaporated on a water-bath until the sweet taste of the sulphinide could no longer be detected. ~~The~~

The solution was allowed to cool and then filtered. The acid was precipitated by means of hydrochloric acid. If the solution is stirred vigorously, after the addition of hydrochloric acid, the acid is precipitated at once as a mass composed of fine crystals. If allowed to stand over night without disturbance large monoclinic crystals are deposited. After filtering, the acid was dissolved in alcohol from which solution it crystallized in large transparent rhombohedrons.

The acid when heated slowly melted at 152°C ; when heated rapidly it melted at 159°C . (uncorrected). It is quite soluble in alcohol and water. Some of the acid was placed in a small boat which was placed in a glass tube with a diameter of about 15 m.m. One end of the tube was connected with a bottle containing a small quantity of sulphuric acid. The other end was connected with a cylinder which contained calcium chloride. By connecting this apparatus with a filter pump, a current of dry air could be drawn through the tube. If a current of air flowing quite rapidly was allowed to pass through the tube, the acid was decomposed when heated to about 128°C . A white sublimate, which was very sweet, collected on the wall of the tube. If the current of air was shut off, the acid decomposed at

135⁰, and white transparent needles of pure benzoic sulphide collected on the wall of the tube. Some of the needles were from ~~ten~~¹⁰ to ~~twelve~~¹² m.m. in length. The acid contains no water of crystallization.

The acid crystallizes in white opaque needles if the solution is allowed to stand at least one week before crystallization takes place. The needles melted at 152⁰. The needles are more soluble in water than the plates. The needle acid contains water of crystallization. Below 110⁰ it lost just one-half a molecule of water.

Calculated for		Found
C_6H_5	COOH	I.
	- 1/2 H ₂ O	4.28
	So ₂ NH ₂	
	H ₂ O 4.29	

The needles when dissolved in alcohol crystallize from the solution in plates which melt at 152⁰.

The needles decomposed when heated to about 115⁰; a white sublimate collected on the walls of the tube. The sublimate was very sweet. Ten grams of benzoic sulphide was added to a caustic soda solution which was prepared by dissolving twenty grams of caustic soda in 250c.c. of water. The solution was treated in the same way as the previous solution. After treating the solution with hydrochloric

acid if it is allowed to stand over night the acid crystallizes out in the form of needles. These proportions give nearly a quantitative yield of the needle acid.

A careful examination of the needles showed that they agreed in every respect with the needles obtained by the recrystallization of the plate acid from water. Melting point 152° ; if the temperature is kept at 152° for a while the acid becomes a solid which melts at 215° . If heated in the apparatus previously described the needles lose one-half molecule of water of crystallization below 100°C .

I - 0.3484 gram of acid lost 0.0150 gram of water at 90° .

II. 0.2920 gram of acid lost 0.0119 gram of water at 98° .

Calculated for	Found.
$\begin{array}{c} \text{C}_6\text{H}_4 \\ \swarrow \text{COOH} \\ \searrow \text{SO}_2\text{NH}_2 \end{array} - 1/2 \text{H}_2\text{O}$	I. II.
H_2O 4.24	4.30 4.11

If a current of air was allowed to flow rapidly through the tube, the acid decomposed at 115°C . and was partially fused. When the current of air was cut off en-

tirely, the acid decomposed at 145° . The acid was heated at this temperature for about six hours. During this time beautiful transparent needles of pure benzoic sulphinide, some of which were several millimeters in length, collected on the walls of the tubes. The needles were extremely sweet and melted at 223° to 224° (uncorrected). By taking proportions intermediate between the proportions used in the previous experiment (fifteen grams of sodium hydroxide with the same amount of benzoic sulphinide ~~acid water~~) a yield is obtained which consists partly of the plate acid and partly of the needle acid. On account of the difference in solubility they can be easily separated, the plates being less soluble than the needles.

20 grams of benzoic sulphinide were added to a caustic soda solution which was prepared by dissolving 20 grams of sodium hydroxide in 750^{cc} water. The solution was treated in the same way as in the previous experiments. From 15 to 20 grams of acid were obtained. The solution was set aside. After standing for a week or two, large transparent, monoclinic plates were deposited, which were fairly soluble in water. The crystals after being dried became translucent. Upon examination the substance was found to be a sodium salt, which, upon analysis, gave the

following result:

- I. 0.1372 gram of the salt gave 0.0347 gram of Na_2SO_4
- II. 0.1805 gram of the salt gave 0.0445 gram of Na_2SO_4
- III. 0.1084 gram of the salt gave 0.0266 gram of Na_2SO_4

Calculated for

Na

Found

I. II. III.

8.16 7.98 7.93

No loss in weight when heated to 165° , neither did it become sweet.

10 grams of benzoic sulphinide were added to a caustic potash solution which was prepared by dissolving 15 grams of potassium hydroxide in 250 c.c. of water. The solution was treated in the same manner as the previous solutions. After treating the solution with hydrochloric acid it was allowed to stand for several days. During this time large monoclinic plates were deposited. The plates became transparent when dried. Upon examination the substance turned out to be a potassium salt which, upon analysis, gave the following results:

- I. 0.2149 gram of salt gave 0.0739 gram of K_2SO_4
- II. 0.1478 gram of salt gave 0.0502 gram of K_2SO_4

Calculated for

K

Found

I. II.

15.40 15.22

No loss in weight when heated to 160° .

20 grams of benzoic sulphinide were added to a caustic soda solution which was prepared by dissolving 30 grams of sodium hydroxide in 500c.c. of water. The solution was treated in the same manner as the other solutions, with the exception of placing the beaker in cold water when the hydrochloric acid was added. When the acid was added to the solution, quite a considerable quantity of heat was evolved. After cooling, a crystalline mass was deposited. This mass when crystallized from water separated as transparent, monoclinic plates which remained transparent after being dried. The substance turned out to be the acid ammonium salt of orthosulphobenzoic acid as evidenced by the fact that ammonia could be detected when the salt was treated with a solution of sodium hydroxide in the cold. Also by the following analysis:

Calculated for		Found.	
		I.	II.
C_6H_4	$COOH$		
	SO_2ONH_4	38.74	39.79
C -	38.35	4.21	4.35
H	4.10	4.65	
N	8.39		

The salt was converted into potassium salt which was analysed:

K	16.25	16.36
---	-------	-------

Salts of Ortho-sulphaminebenzoic Acid.

All of the salts crystallize from a water solution, in well formed crystals. All of them contain water of crystallization with the exception of the mono silver, potassium and ~~common~~ ^{ammonium} salts. Without exception they all become sweet when they lose their water of crystallization; indeed they begin to get sweet just as soon as they begin to lose their water.

All the salts decompose at a few degrees above the temperature at which they give off their water of crystallization.

Sodium Ortho-sulphaminebenzoate -

C_6H_4 $\begin{matrix} \nearrow COONa \\ \searrow SO_2NH_2 \end{matrix}$ + $2\frac{1}{2}H_2O$. This salt was prepared by treating a hot

solution of the acid with pure sodium carbonate to the point of neutralization. The salt crystallizes from an exceedingly concentrated water solution in fine silky needles. Only a few needles were obtained from a solution which was allowed to stand several weeks, in a small test tube. Even then a slight disturbance of the tube would cause the needles to redissolve. As a general rule the salt is deposited as a white amorphous mass. The salt becomes sweet by simply exposing it to the air for a short while. The salt was dried (with drying paper) and analysed at once. The percentage of sodium agreed with $1\frac{1}{2}$ molecules of water, and the percentage of water of crystallization agreed with $2\frac{1}{2}$ molecules of water.

- I. 0.2114 gram of salt lost 0.0357 gram of water below 110° .
- II. 0.2210 gram of salt lost 0.0375 gram of water below 110° .
- III. 0.2628 gram of salt lost 0.0442 gram of water below 110° .

Calculated for		Found		
$\begin{array}{c} \text{COONa} \\ \diagup \\ \text{C}_6\text{H}_4 \\ \diagdown \\ \text{SO}_2\text{NH}_2 \\ \text{H}_2\text{O} \end{array}$	- $2\frac{1}{2}\text{H}_2\text{O}$	I.	II.	III.
	16.79	16.88	16.96	16.81

Calculated for		Found		
		I.	II	III
C_6H_4	$COONa + 1\ 1/2\ H_2O$			
	SO_2NH_2	9.10	9.12	9.11
	Na	9.20		

A few of the needles were obtained and an analysis of the sodium agreed with 2 1/2 molecules of water.

I. 0.0174 grams of salt gave 0.0046 grams of Na_2SO_4 .

II. 0.1945 grams of salt gave 0.0546 grams of Na_2SO_4 .

Calculated for		Found	
		I.	II.
C_6H_4	$COONa - 2\ 1/2\ H_2O$		
	SO_2NH_2	8.62	8.58
	Na	8.58	

Potassium Orthosulphaminebenzoate.

C_6H_4 $\begin{matrix} \nearrow COOK \\ \searrow SO_2NH_2 \end{matrix}$ This salt was prepared by neutralizing the acid with pure potassium carbonate. The salt is extremely soluble in water; Insoluble in methyl alcohol. The salt is deposited from a water solution as a white amorphous mass. It crystallizes from a mixture of alcohol and water in large transparent plates.

I. 0.1445 gram of salt gave 0.0523 gram of K_2SO_4 .

II. 0.1265 gram of salt gave 0.0457 gram of K_2SO_4 .

Calculated for

Found

C_6H_4	COOK	I.	II.
	SO_2NH_2	16.33	16.20
K	16.32		

Calcium Orthosulphamine benzoate.

$$\left(\begin{array}{l} C_6H_4 - COO \\ \quad \quad \quad SO_2NH_2 \end{array} \right)_2 + Ca + 2H_2O \rightarrow$$
 A hot solution of the acid was neutralized with pure calcium carbonate. The salt crystallized from a water solution in well formed translucent crystals.

If crystallization takes place at once, well formed crystals are obtained; but if the solution is allowed to stand several days, the salt is deposited as a white crystalline mass.

I. 0.5475 gram of salt lost 0.0414 gram of water below 140° .

II. 0.3776 gram of salt lost 0.0277 gram of water below 140° .

I. 0.2114 gram of salt gave 0.0598 gram of $CaSO_4$.

II. 0.2926 gram of salt gave 0.0829 gram of $CaSO_4$.

Calculated for

Found

$(C_6H_4 \begin{array}{l} \diagup COO \\ \diagdown SO NH_2 \end{array})_2$	Ca - 2H ₂ O	I	II
		8.27	8.30
Ca	8.40	7.56	7.33
H ₂ O	7.56		

Barium Orthosulphamine benzoate -

$(C_6H_4 \begin{array}{l} \diagup COO \\ \diagdown SO_2NH_2 \end{array})_2 Ba \cdot 3H_2O$ - This salt was prepared by treating the acid with pure barium carbonate. It is quite soluble in water and fairly soluble in alcohol. It crystallizes from a water solution in transparent plates, and sometimes it crystallizes from a water solution in needles in the form of rosettes. It crystallizes in needles from alcohol solution. Indications observed in different experiments, point to the conclusion that the salt crystallizes, under different conditions, with a different number of molecules of water of crystallization. It agrees in every respect with the salt prepared by Mr. W. A. Noyes with the exception of the number of molecules of water of crystallization. Mr. Noyes prepared two salts, one with 2 molecules and another one with 9 molecules of water of crystallization. This salt loses its water of crystallization when


approx. 10. an.

0.1823 gram of the salt lost 0.0171 gram of H_2O below 150° .

I. 0.1214 gram of salt gave 0.0483 gram of $Ba SO_4$

II. 0.1076 gram of salt gave 0.0426 gram of $Ba SO_4$

Calculated for



Found

I II

Ba 23.18

23.39

23.32

H_2O 9.14

9.14

exposed to the air.

Strontium Ortho-sulphaminebenzoate -


$$\left(\text{C}_6\text{H}_4 \begin{array}{l} \text{COO} \\ \text{SO}_2\text{NH}_2 \end{array} \right)_2 \text{Sr} + 2\text{H}_2\text{O} .$$
 This salt is prepared by neutralizing a hot solution of the acid with pure strontium carbonate. If the solution is allowed to evaporate spontaneously the salt crystallizes in long transparent needle-like crystals, some of which are three or four millimeters in length. The crystals remain transparent when exposed to the air.

I. 0.2151 gram of salt lost 0.0150 gram of H_2O below 140° .

I. 0.1385 gram of salt gave 0.0486 gram of Sr SO_4 .

II. 0.2231 gram of salt gave 0.0783 gram of Sr SO_4 .

Calculated for



Found

I II

16.71 16.67

Sr. 16.71

6.97

H_2O 6.87

Zinc Ortho-sulphaminebenzoate -

$$\left(\text{C}_6\text{H}_4 \begin{array}{l} \text{COO} \\ \text{SO}_2\text{NH}_2 \end{array} \right)_2 \text{Zn} - 3\text{H}_2\text{O} .$$
 Two grams of the acid was dis-

solved in a sufficient quantity of water and pure zinc carbonate was added till the solution was neutral. The solution was evaporated to a slight extent, and allowed to stand in a desiccator over night. Short opaque needle-like crystals were deposited during the night. If the solution is allowed to evaporate spontaneously, after several days, large transparent needle-like crystals are deposited, some of which were from one to two centimeters in length. The crystals do not lose their transparency when exposed to the air. The salt is fairly soluble in water.

I. 0.0809 gram of the salt lost 0.0081 gram of water below 130° .

II. 0.0690 gram of salt lost 0.0070 gram of water below 130° .


I. 0.0449 gram of salt gave 0.0137 gram of Zn SO_4 .

II. 0.0621 gram of salt gave 0.0192 gram of Zn SO_4 .

III. 0.1476 gram of salt gave 0.0456 gram of Zn SO_4 .

Calculated for

found.



	I	II	III
	12.24	12.40	12.46

Zn	12.59
H ₂ O	10.39

10.01	10.15
-------	-------

Zinc Ortho-sulphamine benzoate.-

$$\left(\begin{array}{c} \text{COO} \\ \text{C}_6\text{H}_4 \backslash \\ \text{SO}_2\text{NH}_2 \end{array} \right)_2 \text{Zn} + 3\text{H}_2\text{O}.$$
 This salt was prepared from the needle acid in order to compare it with the zinc salt prepared from the plate acid. The two salts resemble one another in every respect with one exception. The solution of this salt, when allowed to evaporate spontaneously deposited large opaque needle-like crystals instead of transparent needles as is the case with the salt prepared from the plate acid.

I. 0.0652 gram of salt gave 0.0201 gram of Zn SO_4 .

I. 0.0138 gram of salt lost 0.0138 gram of water below 150° .

Calculated for		Found
$\left(\begin{array}{c} \text{COO} \\ \text{C}_6\text{H}_4 \backslash \\ \text{SO}_2\text{NH}_2 \end{array} \right)_2$	$\text{Zn} + 3\text{H}_2\text{O}$	I.
Zn	12.59	12.42
H_2O	10.39	10.64

Magnesium Ortho-sulphamine benzoate -

$$\left(\begin{array}{c} \text{COO} \\ \text{C}_6\text{H}_4 \backslash \\ \text{SO}_2\text{NH}_2 \end{array} \right)_2 \text{Mg} + 4\text{H}_2\text{O}.$$
 - This salt was prepared by treating the acid with magnesium hydroxide. The salt

crystallized in well formed transparent needles which are deposited in the form of rosettes. The salt is extremely soluble in water.

I. 0.1850 gram of salt lost 0.0270 below 175°.

II. 0.2458 gram of salt lost 0.0355 gram of water below 175°.


I. 0.2129 gram of salt gave 0.0519 gram of Mg SO_4 .

II. 0.1726 gram of salt gave 0.0425 gram of Mg SO_4 .

III. 0.1416 gram of salt gave 0.0341 gram of Mg SO_4 .

Calculated for

Found



I. II III


4.84 4.92 4.81

Mg 4.84

H_2O 14.52

14.59 14.42

Copper-Ortho-sulphamine benzoate



.- This salt was prepared by boiling a solution of the acid with freshly precipitated copper oxide, in a sand-bath, for about forty minutes. There was deposited after standing over night, large blue feathery needles, some of which were more than a centimeter in length. The salt is insoluble in alcohol and very diffi-

cultly soluble in water. The salt turned a deep green color when it lost its water of crystallization.

I. 0.1369 gram of the salt lost 0.0144 gram of water below 170° .

II. 0.0995 gram of salt lost 0.0095 gram of water below 170°

III. 0.0951 gram of salt lost 0.0085 gram of water below 170° .

I. 0.1684 gram of salt gave 0.0209 gram of Cu O.

II. 0.3018 gram of salt gave 0.0456 gram of Cu O.

Calculated for

Found

$\left(\begin{array}{l} \text{C}_6\text{H}_4 \begin{array}{l} \nearrow \text{COO} \\ \searrow \text{SO}_2\text{NH}_2 \end{array} \end{array} \right) \text{Cu} + 3\text{H}_2\text{O}$		I	II
Cu	12.23	12.41	12.32
H ₂ O	10/44	10.52	10.91

Mono Silver Ortho-sulphamine benzoate. —

This salt was prepared by neutralizing two grams of acid with ammonium hydroxide, and then boiling the solution until no odor of ammonia could be detected. Silver nitrate was added to the solution until no further precipitation

took place. By allowing the solution to stand over night there were deposited, fine feathery needles of a light gray color. The salt is soluble in hot water but insoluble in cold water. Becomes dark ~~colored~~ when exposed to the light.

Although it contains no water of crystallization it becomes sweet when heated up to 150° , and changes its color to light brown.

This salt resembles in every respect the salt prepared by Mr. Noyes (1) by the same method.

I. 0.0875 gram of salt gave 0.0307 gram of Ag.

II. 0.1171 gram of salt gave 0.0412 gram of Ag.

Calculated for

	Found	
	I	II
$\text{C}_6\text{H}_4 \begin{cases} \text{COOAg} \\ \text{SO}_2\text{NH}_2 \end{cases}$	35.08	35.17
Ag	35.06	

Di-Silver Ortho-sulphaminebenzoate,

$\text{C}_6\text{H}_4 \begin{cases} \text{CO Ag} \\ \text{SO}_2\text{NHAg} \end{cases} + \text{H}_2\text{O}.$ - This salt was prepared by neutralizing about two grams of acid with ammonium hydroxide and then adding about $3/4$ as much ammonium hydroxide in excess as was required to neutralize the solution. Silver nitrate

was added until no further precipitation took place. The salt was deposited as a white amorphous mass which was dissolved in boiling hot water. By allowing the solution to stand over night the salt separated out as silver-colored leaflets which became dark ~~colored~~ when exposed to the light. This salt is much less soluble in water than the mono silver salt, and resembles in every respect, with one exception, the di-silver salt prepared by Noyes (1) by treating the barium salt with silver nitrate. This salt contains one molecule of water of crystallization.

The salt was thoroughly dried and was allowed to remain wrapped in drying-paper for about one week. It was then heated to 140° in the apparatus previously described. The current of air was cut off, and water could be distinctly seen as it collected in drops in both ends of the tube. After losing its water it becomes a dark brown color.

I. 0.1231 gram of the salt lost 0.0053 gram of water at 167° .

I. 0.1867 gram of salt gave 0.0933 gram of Ag.

II. 0.1097 gram of salt gave 0.0549 gram of Ag.

(1) Loc. cit.

Calculated for		Found	
		I	II
C_6H_4 \	$COO Ag$	49.97	50.04
	$+ H_2O$		
C_6H_4 \	$SO_2 NHAg$		
	Ag		
	49.68		
H_2O	4.15		


Ammonium Ortho-sulphamine benzoate,

C_6H_4 \ $COO NH_4$. — This was prepared by neutralizing the
 $SO_2 NH_2$
 acid with ammonium hydroxide. The salt is quite soluble
 in water, and crystallizes from a water solution in trans-
 parent plates.

The salt contains no water of crystallization, but
 becomes very sweet when heated to 125°, and decomposes
 quite easily.

Action of Phosphorus Oxychloride on the Sulphamine Acid

Five grams of the acid were mixed with five grams of
 phosphorus oxychloride and heated on a water-bath for two
 hours. No chloride was formed. All of the acid was conver-
 ted into benzoic sulphimide.




~~This reaction is evidence in favor of the formula given to~~

~~the acid by~~ ~~W. R. Noyes.~~ In order to form benzoic sulphide from the acid the hydroxyl group of the carboxyl group has to combine with one hydrogen atom of the amido group of the sulphone group as expressed by the above formula.

Preparation of the Diamide,

$$\text{C}_6\text{H}_4 \begin{cases} \text{CO NH}_2 \\ \text{SO}_2 \text{NH}_2 \end{cases}$$
 The diamide was prepared by the following method:

Ten grams of the acid were mixed with six grams of ammonium cyanate. The mixture was heated, in a closed bath, for three days. During this time it was frequently stirred. As the acid is converted into benzoic sulphinide at quite a low temperature, the mixture was heated from 90° to 100° the first day, from 130° to 135° the second day and from 150° to 160° the third day. The mass, which is of a light straw color, is allowed to cool when it becomes brittle. It is then pulverized and treated with boiling alcohol in which it is soluble. After filtering, the solution is allowed to cool when ~~there~~^{are} ~~is~~ deposited light colored needles. Melting-point 263° .





Analysis of the diamide gave the following results:

- I. 0.1296 gram of diamide gave 0.1528 gram of Ba SO_4 .
- I. 0.1565 gram of the diamide gave 0.0209 gram of nitrogen.
- II. 0.1413 gram of the diamide gave 0.0188 gram of nitrogen.

Calculated for		Found	
C_6H_4	CO NH_2	I.	II.
	SO NH_2	16.20	
S	16.00	13.35	13.27
N	14.00		

I am indebted to Mr. C. E. Waters for the nitrogen determinations. Mr. Waters observed ~~the fact~~ that the ammonia was given off in two different stages. One portion of the ammonia was given off when the sodium hydroxide solution was dilute; after heating sometime the second portion was given off when the soda solution had become very concentrated. ~~So there must be some difference in the stability of the two amide groups.~~

Part II.

ORTHO-CYANAMINEBENZENE-SULPHONIC ACID.

Introduction.

(1)

This acid was first prepared by Fahlberg and B. ^{on the ground,} by the following method: Ortho sulphobenzoic ~~acid~~ when treated with ammonia, reacts according to this equation:



The ammonium salt when treated with silver nitrate is easily converted into the silver salt. The free acid is obtained by treating the silver salt with sulphuric acid.

They describe the acid as crystallizing in thick prismatic crystals which are easily soluble in water and alcohol, and contain one molecule of water of crystallization. Melting-point 185° to 186° , (uncorrected). By melting the acid with potassium hydroxide, ammonia is given off, but when simply heated no ammonia is given off.

(1)

The acid was prepared later by J. ~~as~~ surun by heating orthocyanbenzenesulphonechloride with water or alco-


(1) Ber. d. Chem. Ges. 22, 7598

(2) Ber. d. Chem. Ges. 26, 2288.

hol.

(1)

Mr. Karslake also obtained the acid by hydrolysis of ortho-cyanbenzenesulphonic acid by means of dilute alkali:



Preparation of Acid

The potassium salt of the acid can be obtained in any desired quantity by treating the acid potassium salt of ortho-sulphobenzoic acid with ammonium sulphocyanate (2)

The acid potassium salt was prepared according to (3) the method of Remsen and Saunders.

A modification of the proportions suggested by their method was made for the reason that the benzoic sulphinide used, contained scarcely a trace of para-sulphaminebenzoic acid.

The substances were mixed together in the proportion of five parts of acid potassium salt to three parts of ammonium sulphocyanate. The mixture was heated in a closed

(1) Dissertation, J.H.U., 1896, p.18.

(2) English Patent - Newton & Son - 23, February, 1895.

(3) Dissertation, J.H.U., 1894, p.7.

bath for three days at a temperature from 180° to 200° μ . During this time it was occasionally stirred. The mass which is of a brown color, was allowed to cool when it becomes a stiff cake. It is then finely pulverized and digested with boiling alcohol several times. The potassium salt of the acid which is in the form of sandy crystals was separated from the alcohol by filtering. The salt when recrystallized from water separates out in large transparent plates.

When the potassium salt is treated with silver nitrate, in the proportion of their molecular weights, reaction takes place with the greatest ease with the formation of the silver salt of ortho-carbamine-benzene-sulphonic acid. The acid is obtained by treating the silver salt with hydrochloric acid. The acid crystallizes from a very concentrated water solution in large, thick, transparent crystals which are quite soluble in alcohol. Melting-point 193° - 194° μ (uncorrected). The crystals remain transparent when exposed to the air. It does not become sweet when it loses its one molecule of water of crystallization, neither does it become sweet by continued heating. It decomposes when heated to 125° .

I. 0.2691 gram of the acid lost 0.0228 gram of water below 120° .

II. 0.2585 gram of the acid lost 0.0214 gram of water below 120° .

Calculated for		Found	
$\text{C}_6\text{H}_4 \begin{cases} \nearrow \text{CO NH}_2 \\ \searrow \text{SO}_3\text{OH} \end{cases}$	+ H_2O	I.	II.
		8.47	8.27
	H_2O	8.22	

Salts of Ortho-carbamine-benzenesulphonic acid -

The salts of this acid do not crystallize as well as the salts of ortho-sulphamine-benzoic acid, but are much more soluble in water.

Most of the salts contain water of crystallization but none of them become sweet when they lose their water; neither do they become sweet when they are decomposed.

Potassium ortho-carbamine-benzene-sulphonate 7

$\text{C}_6\text{H}_4 \begin{cases} \nearrow \text{CONH}_2 \\ \searrow \text{SO}_3\text{OK} \end{cases}$ + H_2O .- This salt was prepared by the method described in the preparation of the acid. It crystallizes from a water solution in large transparent

plates. It is quite soluble in water and alcohol. It crystallizes from alcohol in needles. The salt is deliquescent. If the large transparent crystals are placed in a beaker and covered over with water, the moment they become warm a crackling sound can be distinctly heard some distance from the water-bath. The crystals become opaque and crumble to pieces. This phenomenon is doubtless due to the fact that the salt loses its water of crystallization very easily; in this instance ^{probably} it loses it under water.

(1)

A similar salt is described by Karslake ⁽¹⁾ derived from cyanbenzenesulphonic acid, and by Schon ⁽²⁾ derived from the ammonium salt of benzaminesulphonic acid.

I. 0.3439 gram of salt lost 0.0239 gram of water below
100 .

II. 0.5499 gram of salt lost 0.0380 gram of water below
100 .

III. 0.4801 gram of salt lost 0.0331 gram of water below
100 .

I. 0.4660 gram of salt gave 0.1583 gram of K_2SO_4

II. 0.4904 gram of salt gave 0.1657 gram of K_2SO_4 .

III. 0.4246 gram of salt gave 0.1441 gram of K_2SO_4 .

(1) Dissertation, J.H.U., 1895, p.17

" " 1896, p.18

Calculated for			Found		
			I.	II.	III.
C_6H_4	$\begin{array}{l} \nearrow CO NH \\ \searrow SO_3 OK \end{array}$	+ H ₂ O	15.21	15.11	15.21
	K	15.21	6.94	6.91	6.84
	H ₂ O	7.00			

Silver Ortho-carbaminebenzene-sulphonate,

$C_6H_4 \begin{array}{l} \nearrow CO NH_2 \\ \searrow SO_3 OAg \end{array} + H_2O$. — When the potassium salt of the carbamine acid is treated with silver nitrate, in the proportion of their molecular weights, reaction takes place with the greatest ease with the formation of the silver salt. The salt which is fairly soluble in water crystallizes in large, monoclinic, transparent plates. It is decomposed when brought in contact with wet filter paper, and becomes dark ~~colored~~ when exposed to the light for any length of time. The salt is decomposed with the greatest ease when treated with hydrochloric acid or with the chlorides of the metals.

This salt resembles in every respect the salt prepared by Fahlberg and Barge (1) by treating the ammonium salt of the carbamine acid with silver nitrate.

(1) Berichte der D.C. 22, 759.

I. 0.3357 gram of salt lost 0.0193 gram of water below 136 .

II. 0.4136 gram of salt lost 0.0228 gram of water below 136 .

I. 0.2362 gram of salt gave 0.0794 gram of Ag.


II. 0.3681 gram of salt gave 0.1215 gram of Ag.

Calculated for

Found.

		I	II
C_6H_4	$CO NH_2 + H_2O$		
	$SO_2 OAg$	33.19	33.00
Ag	33.14	5.45	5.51
H_2O	5.52		

Tri-Silver Ortho-Carbaminebenzenesulphonate 7



. — A small quantity of the potassium carbaminebenzenesulphonate was dissolved in water, and about two drops of ammonia were added to the solution. A hot solution of silver nitrate was added until no further precipitate was formed. The salt was deposited in a white amorphous mass which was insoluble in boiling water. The salt became dark colored when exposed to the

light and is extremely stable. It was with the greatest difficulty that it was decomposed, therefore, the following analysis cannot be reported with confidence.

Calculated for	Found.
$\text{C}_6\text{H}_4 \begin{cases} \text{CO} \text{ NAg} \\ \text{SO}_3\text{OAg} \end{cases}$	I
Ag 62.07	62.56

Sodium Ortho-Scarbamine-benzene-Sulphonate f

$\text{C}_6\text{H}_4 \begin{cases} \text{CONH}_2 \\ \text{SO}_3\text{ONa} \end{cases} + 2\text{H}_2\text{O}.$ — This salt was prepared by treating the silver salt with sodium chloride.

It is quite soluble in water, from which ~~solution~~ it crystallizes in large, transparent plates which belong to the monoclinic system.

This resembles in every respect the salt described
(1)
by Karslake.

I. 0.1771 gram of the salt lost 0.0244 gram of water below 140° .

II. 0.1572 gram of the salt lost 0.0217 gram of water below 145° .

Calculated for

Found

		I.	II.
$\text{C}_6\text{H}_4 \begin{cases} \text{CONH}_2 \\ \text{SO}_2\text{ONa} \end{cases}$	+ $2\text{H}_2\text{O}$	8.85	8.80
Na	8.98	13.89	13.80
H_2O	13.85		

Barium OrthoCarbamine Benzene Sulphonate

$\text{C}_6\text{H}_4 \begin{cases} \text{CONH}_2 \\ \text{SO}_2\text{O} \end{cases} \text{Ba} + 2\text{H}_2\text{O}$. - This salt was prepared by treating the silver salt with hydrochloric acid, and then adding pure barium carbonate to the point of neutralization. The salt crystallized in transparent plates from a very concentrated water solution.

The percentage of water of crystallization agrees with two molecules of water. The percentage of barium agrees with $2\frac{1}{2}$ molecules of water. The following analyses were made of different portions of the salt prepared at different times.

I. 0.1787 gram of salt lost 0.0114 gram of water below 175° .

II. 0.2194 gram of salt lost 0.0138 gram of water below 175° .

I. 0.0367 gram of salt gave 0.0357 gram of Ba SO₄.

II. 0.1584 gram of salt gave 0.0537 gram of Ba SO₄.

Calculated for

Found.

$\left(\text{C}_6\text{H}_4 \begin{array}{l} \diagup \text{CO NH}_2 \\ \diagdown \text{SO}_2\text{O} \end{array} \right)_2$	Ba + 2H ₂ O		Found.	
			I	II
			6.37	6.28
H ₂ O	6.38			

Calculated for

Found.

$\left(\text{C}_6\text{H}_4 \begin{array}{l} \diagup \text{CO NH}_2 \\ \diagdown \text{SO}_2\text{O} \end{array} \right)_2$	Ba + 2 1/2 H ₂ O		Found.	
			I	II
			23.56	23.61
Ba	23.54			

Strontium Ortho-carbaminebenzenesulphonate ;

$\left(\text{C}_6\text{H}_4 \begin{array}{l} \diagup \text{CO NH}_2 \\ \diagdown \text{SO}_2\text{O} \end{array} \right)_2$ This salt was prepared just as the corresponding barium salt by using strontium carbonate. The salt separated, out of an extremely concentrated water solution, as small, transparent crystals with no definite form.

I. 0.2145 gram of the salt gave 0.0815 gram of Sr SO₄.

II. 0.2017 gram of the salt gave 0.0764 gram of Sr SO₄.

Calculated for

Found

$\left(\text{C}_6\text{H}_4 \begin{array}{l} \diagup \text{CO NH}_2 \\ \diagdown \text{SO}_2\text{O} \end{array} \right)_2$	Sr.		Found	
			18.08	18.04
Sr	17.85			

Calcium Ortho-carbamine-benzene-sulphonate 7

$$\left(\begin{array}{l} \text{C}_6\text{H}_5 \text{---} \text{CONH}_2 \\ \text{SO}_3\text{O} \end{array} \right) \text{Ca} + 2\text{H}_2\text{O} \text{---}$$
 This salt was prepared just as the corresponding strontium salt by using pure calcium carbonate. The salt crystallized in large transparent plates from a very concentrated water solution.

I. 0.1333 gram of salt lost 0.0097 gram of water below 185°.

II. 0.1986 gram of salt lost 0.0150 gram of water below 185°.

I. 0.1169 gram of salt gave 0.0332 gram of CaSO_4 .

II. 0.1127 gram of salt gave 0.0323 gram of CaSO_4 .

Calculated for

Found

$\left(\begin{array}{l} \text{C}_6\text{H}_5 \text{---} \text{CONH}_2 \\ \text{SO}_3\text{O} \end{array} \right) \text{Ca} + 2\text{H}_2\text{O}$		
	I	II
	7.28	7.35
Ca 7.56	8.39	8.43
H ₂ O 8.40		

Zinc Ortho-carbamine-benzene-sulphonate 7

$$\left(\begin{array}{l} \text{C}_6\text{H}_5 \text{---} \text{CONH}_2 \\ \text{SO}_3\text{O} \end{array} \right) \text{Zn} + 3\text{H}_2\text{O} \text{---}$$
 This salt was prepared by treating the acid with pure zinc carbonate. The salt is

Deposited from a very concentrated water solution in the form of short thick needles. It is efflorescent.

I. 0.2187 gram of salt lost 0.0227 gram of water below 150°.

II. 0.2477 gram of salt lost 0.0256 gram of H₂O below 150°

I. 0.0934 gram of salt gave 0.0286 gram of ZnSO₄.

II. 0.1022 gram of salt gave 0.0320 gram of ZnSO₄.

Calculated for		Found	
		I	II
$\left(\begin{array}{c} \text{CONH}_2 \\ \text{C}_6\text{H}_4 \backslash \\ \text{SO}_2\text{O} \end{array} \right) \text{Zn} + 3\text{H}_2\text{O}$			
		12.31	12.62
Zn	12.59	10.37	10.34
H ₂ O	10.39		

Magnesium carbamine benzene sulphonate ,

$$\left(\begin{array}{c} \text{CONH}_2 \\ \text{C}_6\text{H}_4 \backslash \\ \text{SO}_2\text{O} \end{array} \right) \text{Mg} + 2\text{H}_2\text{O}.$$
 This salt was prepared just as the zinc salt by using pure magnesium carbonate. The salt separates out as a crystalline mass from a very concentrated water solution.

I. 0.1983 gram of salt lost 0.0146 gram of water below 140° .

II. 0.2002 gram of salt lost 0.0158 gram of water below 140° .

I. 0.1204 gram of salt gave 0.0310 gram of MgSO_4 .

II. 0.1904 gram of salt gave 0.0490 gram of MgSO_4 .

Calculated for

Found

$(\text{C}_6\text{H}_4 \begin{array}{l} \text{CONH}_2 \\ \text{SO}_2\text{O} \end{array}) \text{Mg} + 2\text{H}_2\text{O}$		I	II
		5.14	5.14
		7.36	7.39
Mg	5.22		
H_2O	7.82		

Copper Ortho-carbamino-benzene-sulphonate ,

$(\text{C}_6\text{H}_4 \begin{array}{l} \text{CONH}_2 \\ \text{SO}_2\text{O} \end{array}) \text{Cu} + 4\text{H}_2\text{O}$. — This salt was prepared by boiling the acid with copper oxide on a water-bath. When allowed to cool, the salt was deposited as beautiful deep blue plates. When the salt loses its water of crystallization it becomes gray ~~colored~~. If a small quantity of water is added to the gray ~~colored~~ salt it becomes light blue ~~colored~~.

I. 0.1761 gram of salt lost 0.0240 gram of water below 200°.

II. 0.1760 gram of salt lost 0.0240 gram of water below 200°.

I. 0.1083 gram of salt gave 0.0127 gram of CuO.

II. 0.0435 gram of salt gave 0.0064 gram of CuO.

Calculated for		Found	
$\begin{array}{l} \text{CONH}_2 \\ \diagup \\ \text{C}_6\text{H}_4 \\ \diagdown \\ \text{SO}_2\text{O} \end{array} \quad \begin{array}{l}) \\) \\) \end{array} \quad \text{Cu} + 4\text{H}_2\text{O},$		I	I
		11.72	11.72
Cu	11.86	13.62	13.63
H ₂ O	13.45		

Ammonium Ortho-carbaminebenzene-sulphonate ,

$$\begin{array}{l} \text{CONH}_2 \\ \diagup \\ \text{C}_6\text{H}_4 \\ \diagdown \\ \text{SO}_2\text{ONH}_4 \end{array} \quad , -$$
 This salt was prepared by neutralizing the acid with ammonium hydroxide. It crystallized from a concentrated water solution in large plates. One plate obtained was at least 2 1/2 c.c. in length. It gives off ammonia when treated with sodium hydroxide in the cold. After heating the salt up to 140° no sweet taste could be

detected. Melts at 262° to 263° (uncorrected). If the current of air was cut off water collected in the ends of the tube when the salt was heated to 125° .

I. 0.1503 gram of salt lost 0.0062 gram of water below 125° .

II. 0.1404 gram of salt lost 0.0059 gram of water below 125° .

I. 0.1562 gram of salt gave 0.1649 gram of Ba SO_4 .

Calculated for		Found	
$\begin{array}{c} \text{CONH}_2 \\ \diagup \\ \text{C}_6\text{H}_4 \\ \diagdown \\ \text{SO}_2\text{ONH}_4 \end{array}$	+ $1/2 \text{ H}_2\text{O}$	I.	II
		4.12	4.20
H_2O	3.96		

Calculated for		Found
$\begin{array}{c} \text{CONH}_2 \\ \diagup \\ \text{C}_6\text{H}_4 \\ \diagdown \\ \text{SO}_2\text{ONH}_4 \end{array}$		I
		14.46
S.	14.67	

(1)

This salt was prepared by Fahlberg and Barge and described as crystallizing from alcohol in white needles which melt at 225° - 226° . (2) Jepsurun describes it as crystal-

(1) Loc. cit.

(2) Loc. cit.

(3) Loc. cit.

lizing from water in granular crystals containing one molecule of water of crystallization. Karslake⁽¹⁾ describes the salt as crystallizing from a water solution, in transparent needles, which contain no water of crystallization and which melted at $219^{\circ} - 220^{\circ}$ (uncorrected).

Preparation of Chloride.

Five grams of the potassium salt was mixed with five grams of phosphorus oxychloride in a small flask. The flask was attached to a return condenser, and then placed in a bath which contained boiling water. The mixture was heated for about one-half an hour. The white mass was pulverized in a mortar and treated with cold water, the ortho-carbamine-benzene-sulphon~~e~~chloride, which remains in the form of white crystalline powder insoluble in water, is filtered off. It was then thoroughly dried. It crystallized from chloroform solution in small thick opaque plates.

Melting-point 63° .




I. 0.1156 gram of salt gave 0.0984 gram of AgCl (Carion² method).

(1) Loc. cit.

Calculated for

Found




I.
16.00

Cl 16.17

Potassium Orthocyanbenzenesulphonate

$\begin{array}{l} \text{CN} \\ \text{C}_6\text{H}_4 \backslash \text{SO}_2\text{OK} \end{array}$. - This salt was prepared by treating the potassium salt with phosphorus oxychloride as in the previous experiment, only the mixture in this experiment was simply warmed in the water-bath. The whole of the potassium salt was converted into the cyan salt. It crystallizes from a moderately dilute water solution in small transparent needles radiating from a common centre.

If the mixture of the potassium salt and phosphorus oxychloride is placed in a cold water bath, and then heated to the boiling temperature, the potassium salt will be partly converted into the chloride and partly into the cyan potassium salt.



~~This reaction is evidence in favor of the formula given to the acid by Fahlberg and Bange.~~

(2) This salt resembles in every respect the salt described by Noyes, (1) derived from ammonium cyanbenzenesulphonate.

Calculated for

Found.

$\begin{array}{c} \text{C}_6\text{H}_4 \\ \text{K} \end{array}$	CN		
	SO ₂ OK		
	17.67	I	II
		17.67	17.71


A small portion of the chloride was placed in a beaker which was then placed in a dish of cold water. The chloride was treated with a small quantity of an ordinary solution of ammonium hydroxide. The beaker was allowed to stand in the water for about three hours, being stirred at frequent intervals. It was then warmed on a water-bath for about half an hour. The solution was then filtered and a white crystalline substance remained on the filter paper. It was soluble only in boiling alcohol from which solution it crystallized in small opaque monoclinic (?) plates. The substance did not melt when heated to the boiling-point of sulphuric acid. A sulphur determination was made by the Liebig method. Ammonia was given off when the substance was fused with potassium hydroxide, although not as freely as in the case of the sulphamine diamide. This substance seemed to be more stable than the sulphamine diamide.

I (1885-1886)

I. 0.1203 gram of substance gave 0.1506 gram of Ba SO₄.

Calculated for

Found



I

17.20

Possibly the substance was not thoroughly oxidized when fused with potassium hydroxide or else it is not the carbamine diamide.

Appendix.

The following work was done by Mr. Benjamin Palmer Caldwell, a student in the department of Physical Chemistry -

Determination of the Molecular Conductivity and Activity Coefficient (Dissociation Constant) of Ortho-Sulphamine Benzoic Acid, C_6H_4 $\begin{matrix} \nearrow \text{COOH} \\ \searrow \text{SO}_2\text{NH}_2 \end{matrix}$ and Ortho-Carbamine Benzoic Acid, C_6H_4 $\begin{matrix} \nearrow \text{CONH}_2 \\ \searrow \text{SO}_2\text{OH} \end{matrix}$ *benzene sulphonic*

The conductivity was determined by the method of Kohlrausch (Zeit. phys. Chem., I, 105; 2, 902) using a Wheatstone bridge, induction coil, and telephone.

Since organic acids, in general, are not strongly dissociated in solution, complete dissociation is only reached at dilutions far too great to permit the use of the conductivity method. The conductivity of the sodium salt at infinite dilution is, therefore, ^{assumed} ~~obtained~~, since this is strongly dissociated, and that of the acid is computed from this value by subtracting the constant for the Sodium ion, 49, and adding to the remainder the constant for the Hydrogen ion, 325. In the case of the Ortho-sulphamine Benzoic acid, this was not done, since the composition of the dry

sodium salt was not positively ascertained. The value 360, was taken as the molecular conductivity of this acid at complete dissociation, it having been seen in the examination of the results of Ostwald's work on organic acids (Zeit. Phys. Chem. 3, 170, 240, 360.) that the value for a very large number of closely related monobasic acids fell between 357 and 362. This number, for monobasic acids is the sum of the ^{free} Hydrogen constant, 325, and the constant for the union made up of the rest of the acid. This latter constant is nearly the same throughout, for a large number of acids. The value, 360, cannot be wrong by more than two units - which would not ~~materially~~ ^{materially} affect the results dependent upon it ~~materially~~.

In the case of the Ortho-carbamine ^{benzene-sulphonic} ~~Sulpho-Benzotic~~ acid, the sodium salt was employed, and the value for the molecular conductivity of that acid at complete dissociation was computed as shown above.

The amount of dissociation at any specified dilution, α , is obtained by dividing the molecular conductivity at that dilution μ_v , by the molecular conductivity at infinite dilution μ_∞ - that is, at complete dissociation: $\alpha = \frac{\mu_v}{\mu_\infty}$.

The chemical activity of acids is directly propor-

tional to the dissociation constant, k . This is obtained by the formula of Ostwald (Zeit. Phys. Chem., 3, 172), which was ~~deduced~~^{deduced} from the condition of a dissociating vapor

$$A_2 = \frac{cA^2}{(1-A)V}$$

Where \underline{V} is the number of litres of solution containing a gram molecule of the dissolved substance. This formula holds for weakly dissociated substances, but fails when applied to strongly dissociated compounds.

The results obtained by Mr. Caldwell are given below with the acids, submitted ~~to~~^{by} me for examination.

COOH C ₆ H ₄ \ SO ₂ NH ₂		Molec. Wt. 201.146	Temperature 25°
Volume	Conduc- tivity,	Dissociation, 100	Dissociation con- stant, 100 k.
25	75.95	21.09	0.2254
50	101.74	28.23	0.2221
100	133.89	37.19	0.2361
200	171.55	47.65	0.2264
400	212.48	59.02	0.2125
800	251.10	69.07	0.1928
1600	278.12	77.02	-
3200	300.14	83.37	-
6400	323.41	89.83	-

12800 334.21 92.83 -

$\Lambda_{\infty} = 360$ 100k (mean of first five) 0.2245.

I omit the last four values for the dissociation constant. The conductivity of the water at high dilutions brings in a slight error in the molecular conductivity as determined, and slight errors there, are often magnified ten times or more in the calculation of dissociation constants.

CONH₂
C₆H₅SO₃ONa + 2H₂O Molec. Wt. 259 213. Temperature, 25°

Volume	Conductivity
20	69.056
40	72.805
80	75.910
160	78.530
320	80.610
640	82.352
1280	85.506
2560	87.140

The error from the water commences to show itself at N/1280 dilution. The correct conductivity for complete dissociation is about 85.

85 - 49 + 325 = 361, μ_{H} of the acid.

$\text{C}_6\text{H}_4 \begin{array}{l} \text{CONH} \\ \diagdown \\ \text{SO OH} \end{array} + \text{H}_2\text{O} \quad \text{Molec. wt. 219} \quad 152.$
 $\mu_{\text{H}} = 361. \quad \text{Temperature } 25$

Volume.	Conductivity,	Dissociation, 100
50	317.01	87.81
100	326.53	90.45
200	331.96	91.95
400	334.65	92.70
800	334.68	92.70
1600	336.52	93.21
3200	344.70	95.48
6400	357.26	99.02
12800	Too dilute to work.	

100 k = 12.64.

The water used in all of the above work was distilled from acid potassium bichromate and alkaline potassium permanganate in the way described by Jones and Mackay in (Am.Chem.Jour., 19,91). Zitsche Phys. Chem. 32,237.

~~Benjamin Palmer Caldwell.~~

Johns Hopkins University,
Baltimore, Maryland.

Biographical.

Francis Daniel Wilson was born in Chatham, Virginia, December 3, 1871. His primary education was received in the public school of that town. After engaging in the mercantile business for three years he entered the Virginia Polytechnic Institute at Blacksburg, Virginia in fall of 1890, and received the degree of Bachelor of Science in 1894, and Master of Science in 1896. During the two years, from 1894 to 1896, he was assistant in General and Analytical Chemistry. In the fall of 1896 he entered the Johns Hopkins as a student in Chemistry, Geology and Mineralogy.

Each year he was appointed to ordinary Virginia scholarships.









